

Linovera®

Rethink skin health

Hyperoxygenated fatty acids
(HOFA) & vegetable extracts

B|BRAUN
SHARING EXPERTISE




Rethink skin health


Chronic wounds are a major medical concern world-wide. They significantly impact patients' quality of life, while also affecting close family and relatives.^{1,2,3} In addition, chronic wounds place a considerable financial burden on healthcare systems and society.⁴

Topical preventive agents such as Linovera® can help to preserve patients' skin health and contribute to overall cost reductions. They can be used to enhance skin integrity as they maintain skin moisture levels and prevent dehydration.

Linovera® ingredients


The hydroxygenated fatty acids (HOFA) in Linovera® are derived from sunflower oil, which is naturally rich in linoleic acid – an essential fatty acid that is important for maintaining epidermal integrity and the skin's natural water barrier.^{5,6}





Mode of action

Linovera[®] can be useful in the prevention of skin breakdown or in the management of already impaired skin.^{10,12}



*TEWL: Transepidermal Water Loss

How to use Linovera®


- 1 Clean the skin at risk or the periwound skin.
- 2 Apply the product and spread it gently over the skin, rather than rubbing it in.
- 3 The product should be applied 2 or 3 times a day.

Protective dressings

If necessary, protect risk areas with dressings suitable for each location.



Easy to use and apply on large surfaces



Ordering information



LINOVERA®

Product	B. Braun code	Quantity
Linovera® 30 ml	450080	1
Linovera® 30 ml	450082	10
Linovera® 50 ml	450081	10



LINOVERA® EMULSION

Product	B. Braun code	Quantity
Linovera® emulsion 50 ml	451078	1
Linovera® emulsion 100 ml	451076	1

References

- 1 Garraud O, Hozzein WN, Badr G. Wound healing: time to look for intelligent, natural 'immunological approaches? BMC Immunol. 2017 Jun 21;18(Suppl 1):23. doi: 10.1186/s12865-017-0207-y. PMID: 28681702; PMCID: PMC5499069.
- 2 Vogt TN, Koller FJ, Santos PND, Lenhani BE, Guimarães PRB, Kalinke LP. Quality of life assessment in chronic wound patients using the Wound-QoL and FLQA-Wk instruments. Invest Educ Enferm. 2020 Oct;38(3): e11. doi: 10.17533/udea.iee.v38n3e11. PMID: 33306901; PMCID: PMC7885545.
- 3 Probst S, Gschwind G, Murphy L, Sezgin D, Carr P, McIntosh C, Gethin G. Patients' acceptance' of chronic wound-associated pain – A qualitative descriptive study, Journal of Tissue Viability, Volume 32, Issue 4, 2023, Pages 455-459, ISSN 0965-206X.
- 4 Nussbaum SR, Carter MJ, Fife CE, DaVanzo J, Haught R, Nussgart M, Cartwright D. An Economic Evaluation of the Impact, Cost, and Medicare Policy Implications of Chronic Nonhealing Wounds. Value Health. 2018 Jan;21(1):27-32. doi: 10.1016/j.jval.2017.07.007. Epub 2017 Sep 19. PMID: 29304937.
- 5 Lin TK, Zhong L, Santiago JL. Anti-Inflammatory and Skin Barrier Repair Effects of Topical application of Some Plant Oils. Int J Mol Sci. 2017 ec 27;19(1):70. doi: 10.3390/ijms19010070. PMID: 9280987; PMCID: PMC5796020.
- 6 Lania BG, Morari J, Almeida ARd, Silva MNd, Vieira-Damiani G, Lins KdA, César CL, Velloso LA, Maia NB, Cintra ML, Velho PENF. Topical essential fatty acid oil on wounds: Local and systemic effects. PLoS One. 2019;14(1):e0210059.
- 7 Aloweni F, Lim M, Chua T, Tan S, Lian S, Ang S. A randomised controlled trial to evaluate the incremental effectiveness of a prophylactic dressing and fatty acids oil in the prevention of pressure injuries. Wound Practice & Research: Journal of the Australian Wound Management Association. 2017;25(1):24-34.
- 8 Martín BC, Martín JIC, Oliver JL, Gómez JD. The Effect of Hyperoxygenated Fatty Acids in Preventing Skin Lesions Caused by Surgical Pneumatic Tourniquets. Adv Skin Wound Care. 2018 May;31(5): 214-217. doi: 10.1097/01.ASW.0000530065.52255.5d. PMID: 29672392.
- 9 Otero DP, Domínguez DV, Fernández LH, Magariño AS, González VJ, Klepzing JVG, Montesinos JVB. Preventing facial pressure ulcers in patients under non-invasive mechanical ventilation: a randomised control trial. J Wound Care. 2017;26(3):128-36.
- 10 Federici A, Federici G. Evaluation of the efficacy of a preparation containing essential fatty acids and plant extracts in the prevention and treatment of dehydrated skin in diabetic patients. Acta Vulnologica. 2014;12(4):177-86.
- 11 Linovera & Linovera Emulsion-PMS_09.AN2. Questionnaire Report: Informe de cuestionario de valoración de producto [Report on the product rating questionnaire]. Rubí (Barcelona), Spain: Braun Medical S.A.U. 2021. In Spanish.
- 12 Torra i Bou JE, Segovia Gómez T, Verdú Soriano J, Nolasco Bonmatí A, Rueda López J, Arboix i Perejamo M. The effectiveness of a hyperoxygenated fatty acid compound in preventing pressure ulcers. J Wound Care. 2005;14(3):117-21.